



## OLIGO-1,6-GLUCOSIDASE (microbial) (Lot 120501b)

### Recombinant

#### E-OAGUM

02/16

(EC 3.2.1.10) oligosaccharide 6- $\alpha$ -glucohydrolase

CAZy: GH Family 13

#### PROPERTIES

##### 1. ELECTROPHORETIC PURITY

- Single band on SDS-gel electrophoresis (MW ~ 67,300)
- Single major band on isoelectric focusing (pI ~ 5.4)

##### 2. SPECIFIC ACTIVITY

**171 U/mg protein (on *p*-nitrophenyl- $\alpha$ -D-glucopyranoside) at pH 4.5 and 40°C.**

~ 321 U/mg protein (on *p*-nitrophenyl- $\alpha$ -D-glucopyranoside) at pH 4.5 and 50°C;

**One Unit** of oligo-1,6-glucosidase activity is defined as the amount of enzyme required to release one  $\mu$ mole of *p*-nitrophenol (*p*-NP) per minute from *p*-nitrophenyl- $\alpha$ -D-glucopyranoside (5 mM) in sodium acetate buffer (50 mM), pH 4.5.

##### 3. RELATIVE RATES OF HYDROLYSIS OF SUBSTRATES:

Substrate	%
<i>p</i> -NP- $\alpha$ -D-Glucopyranoside	100
<i>p</i> -NP- $\beta$ -D-Glucopyranoside	< 0.001
<i>p</i> -NP- $\alpha$ -D-Galactopyranoside	< 0.001
<i>p</i> -NP- $\beta$ -D-Galactopyranoside	< 0.001
<i>p</i> -NP- $\alpha$ -D-Mannopyranoside	~ 0.05
<i>p</i> -NP- $\beta$ -D-Mannopyranoside	< 0.001
<i>p</i> -NP- $\alpha$ -D-Xylopyranoside	< 0.001
<i>p</i> -NP- $\beta$ -D-Xylopyranoside	< 0.001
6 <sup>3</sup> - $\alpha$ -D-Glucosyl-maltotriose (O-GMT)	~ 13
6 <sup>3</sup> - $\alpha$ -D-Glucosyl-maltotriosyl-maltotriose (O-GMH)	~ 0.57
Isomaltose	~ 11.5
Lactose	< 0.001
Maltose	< 0.01
Panose	~ 21
Sucrose	~ 0.29
Trehalose	< 0.01

Action on disaccharide and *p*-NP substrates was determined at final concentration of 5 mg/mL and 5 mM respectively, in sodium acetate buffer (100 mM), pH 4.5 at 40°C.

##### 4. PHYSICOCHEMICAL PROPERTIES:

Recommended conditions of use are at pH 4.5 and 40°C - 50°C.

pH Optima: 4.5

pH Stability: 5.0 - 9.0 (> 75% control activity after 24 hours at 4°C)

Temperature Optima: 50°C (10 min. reaction)

Temperature Stability: up to 40°C

**5. STORAGE CONDITIONS**

The enzyme is supplied as an ammonium sulphate suspension in 0.02% (w/v) sodium azide and should be stored at 4°C. For assay, this enzyme should be diluted in sodium acetate buffer (50 mM), pH 4.5 containing 1 mg/mL BSA. **Swirl to mix the enzyme immediately prior to use.**