

ARABINAZYME

Tablets

ASSAY PROTOCOL

T-ARZ-200T

02/22

200 Tablets

 [Play Training Video](#)



Megazyme

SUBSTRATE:

The substrate employed is Azurine-crosslinked-debranched arabinan (AZCL-Arabinan). Highly purified arabinan from sugar-beet pulp is treated with α -L-arabinofuranosidase to remove 1,3- and 1,2- α -linked arabinofuranosyl residues, leaving linear 1,5- α -L-arabinan. This polysaccharide still contains a small percentage of galacturonic acid, galactose and rhamnose (6, 4 and 2%, respectively), but is resistant to attack by polygalacturonanase and *endo*-1,4- β -D-galactanase. The polysaccharide is then dyed and crosslinked. Treatment of this substrate with a large excess of α -L-arabinofuranosidase results in a limited release of arabinose but no release of dye labelled fragments.

AZCL-Arabinan is a highly sensitive and very specific substrate for the assay of *endo*-arabinanase and can be used to measure this activity in the presence of large excesses of other pectin degrading enzymes and in the presence of sugars and other stabilisers present in industrial powder and liquid pectinase enzyme preparations. This substrate is supplied commercially in a ready-to-use form as **Arabinazyme tablets**.

EXTRACTION/DILUTION BUFFER:

Sodium Acetate buffer (50 mM, pH 4.0) containing sodium azide (0.02%)

Glacial acetic acid (3.05 g, 1.05 g/mL) is added to 900 mL of distilled water. This solution is adjusted to pH 4.0 by the addition of 1 M (4 g/100 mL) sodium hydroxide solution. Approximately 12 mL is required.

Sodium azide (0.2 g) is added, the pH is readjusted to pH 4.0 by dropwise addition of 1 M hydrochloric acid, and the volume adjusted to 1 litre. Store at 4°C.

NOTE: Sodium azide is a toxic chemical and should be treated accordingly.

ENZYME DILUTION:

For most commercial pectinase enzyme preparations, a dilution of 500-fold is required. Liquid enzyme preparation (1.0 mL) is added (using a positive displacement dispenser) to **Extraction/Dilution** buffer (9.0 mL). An aliquot of this solution (0.2 mL) is then further diluted by addition to 9.8 mL of sodium acetate buffer (50 mM, pH 4.0).

With powder samples, the preparation (1.0 g) is added to **Extraction/Dilution** buffer (20 mL) and mixed until either completely dissolved or dispersed. This solution is clarified by centrifugation (1,000 g, 10 min) or filtration through Whatman No.1 filter circles, and then further diluted as for the liquid samples.

If reaction values greater than 1.5 **Absorbance Units** are obtained, the enzyme solution should be further diluted in Extraction/Dilution buffer and the assay repeated.

ASSAY PROCEDURE:

1. Aliquots (0.5 mL) of suitably diluted enzyme in Extraction/Dilution buffer are pre-equilibrated to 40°C for 5 min.
2. Reaction is initiated by the addition of an Arabinazyme tablet. The tablet hydrates rapidly. The suspension should **not** be stirred.
3. After exactly 10 min at 40°C, the reaction is terminated by the addition of Trizma Base solution (10 mL, 2% w/v) with vigorous stirring on a vortex mixer.
4. After about 5 min standing at room temperature, the slurry is stirred again and filtered through a Whatman No. 1 (9 cm) filter circle.
5. The absorbance of the reaction solutions are then measured at 590 nm against the reaction blank.

A **substrate/enzyme (reaction) blank** is prepared by adding Trizma Base (10 mL) to the enzyme solution (0.5 mL) before the addition of the Arabinazyme tablet.

A **single blank** is required for each set of determinations and this is used to zero the spectrophotometer.

STANDARDISATION:

A standard curve relating the activity of purified **endo-arabinanase** from *Aspergillus niger* on linear 1,5- α -L-arabinan [carboxymethyl form] and Arabinazyme (Lot 111201) is shown in Figure 1. Enzyme activity on CM-linear 1,5- α -L-arabinan was determined at a substrate concentration of 2 mg/mL in 100 mM sodium acetate buffer (pH 4.0) at 40°C using the Nelson/Somogyi reducing sugar procedure. The effects of pH and salt concentrations on measured activity are shown in Figures 2 and 3.

One **Unit of activity** is defined as the amount of enzyme required to release one micromole of arabinose reducing-sugar equivalents from

CM-linear arabinan per minute under the defined assay conditions. The rate of hydrolysis of CM-linear arabinan by *endo*-arabinanase is the same as that obtained for linear arabinan recovered from pear juice concentrate.

CALCULATION OF ACTIVITY:

endo-Arabinanase activity in the sample being assayed is determined by reference to the standard curve or to a regression equation for Arabinazyme tablets (Figure 1).

The regression equation varies slightly from batch to batch of Arabinazyme tablets. To minimise this inconvenience, large batches of tablets are prepared.

For Arabinazyme Lot 111201:

endo-Arabinanase milli-Units/assay (i.e. 0.5 mL)

$$= 14.5 \times \text{Abs.}^2 + 39.3 \times \text{Abs.}$$

endo-Arabinanase activity per mL or g of original preparation:

$$= \text{milli-Units/assay} \times 2 \times \frac{1}{1000} \times \text{Dilution}$$

where:

2 = conversion from 0.5 mL (as assayed) to 1.0 mL.

1/1000 = conversion from milliUnits to Units.

Dilution = total dilution of the original enzyme preparation.

$$\text{mU/assay} = 14.5 \times \text{Abs.}^2 + 39.3 \times \text{Abs.}$$




Figure 1. *endo*-Arabinanase (*A. niger*) standard curve on Arabinazyme tablets (Lot 111201).




Figure 2. Effect of pH on the activity of *endo*-arabinanase on Arabinazyme tablets.




Figure 3. Effect of buffer salt concentration on the activity of *endo*-arabinanase on Arabinazyme tablets



Megazyme

North America

NEOGEN Headquarters
800.234.5333 (USA/Canada)
foodsafety@neogen.com
NEOGEN.com

Europe, Middle East and Africa

NEOGEN Europe
+ 44 (0) 1292 525 600
info_uk@neogeneurope.com
NEOGEN.com

Mexico

NEOGEN Latinoamérica
+52 (55) 5254-8235
informacion@neogenlac.com
NEOGEN.com

Brazil

NEOGEN do Brasil
+55 19 3935.3727
info@neogendobrasil.com.br
NEOGEN.com

China

NEOGEN Bio-Scientific Technology
+86 21 6271 7013
info@neogenchina.com.cn
www.neogenchina.com.cn

India

NEOGEN Food and Animal Security
+91 484 2306598, 2301582
info@neogenindia.com
www.neogenindia.com

NEOGEN

Megazyme Building 1,
Bray Business Park, Bray,
Co. Wicklow,
A98 YV29
IRELAND.
+353 (1) 286 1220
Megazyme.com

WITHOUT GUARANTEE

The information contained in this assay protocol is, to the best of our knowledge, true and accurate, but since the conditions of use are beyond our control, no warranty is given or is implied in respect of any recommendation or suggestions which may be made or that any use will not infringe any patents.